



FMDH
PUBLIC HEALTH

📍 621 3rd Street South
Southwing Entrance
Glasgow, MT 59230
☎ (406) 228-3616
📠 (406) 228-3509
🌐 fmdh.org/publichealth
📘 FMDH Public Health
📷 @fmdhpublichealth

Immunization Schedule

To schedule an appointment, call (406) 228-3616

At Birth	Hep B							
2 months	D TaP	Hep B	Polio	Hib	Pcv20	RVV		
4 months	D TaP		Polio	Hib	Pcv20	RVV		
6 months	D TaP	Hep B	Polio		Pcv20	RVV		
12 months		Hep A		Hib	Pcv20		MMR	
15 months	D TaP					VZV		
18 months		Hep A						
4 - 6 years old	D TaP		Polio			VZV	MMR	
11 - 12 years old	Tdap			MCV4			HPV9	HPV9
16 - 17 years old				MCV4				
Adults/Seasonal	Tdap	Flu	Covid		RSV			

- DTaP - Diphtheria, Tetanus, (acellular) Pertussis
- IPV - Injectable Polio Vaccine
- Hep B - HBV - Hepatitis B Vaccine
- Hep A - HAV - Hepatitis A Vaccine
- Hib - Haemophilus Influenza type B
- MMR - Measles, Mumps, Rubella
- VZV - Varicella (Chickenpox)
- PCV13 - Pneumococcal Conjugate Vaccine
- RVV - Rotavirus Vaccine

DTaP/Polio/Hep B = combination vaccine called Pediarix

Infant cannot receive this vaccine if the first dose is not given prior to age 15 weeks.

Vaccines for 4 year olds:

- DTaP
- IPV
- MMR
- VZV


WHAT ARE NORMAL VACCINE INJECTION REACTIONS?



Pain & Swelling at Injection Site

WHAT TO DO:

- Movement
- Ice pack
- If skin is red and swollen, apply hydrocortisone cream



Fatigue

WHAT TO DO:

- Rest as needed



Fever, Chills or Headache

WHAT TO DO:

- Tylenol or Ibuprofen
- Rest
- Drink plenty of fluids

Muscle Aches & Soreness



WHAT TO DO:


- Tylenol or Ibuprofen
- Rest
- Heat or Ice pack

WHEN SHOULD I SEE MY DOCTOR AT THE CLINIC?



Fever that lingers longer than 3-4 days

Pain and swelling have spread beyond the injection site after 3-4 days. Localized bump may linger for a week.



SIDE EFFECTS ARE A SIGN THAT THE VACCINE IS WORKING.

Mild to moderate reactions to vaccines are common and are a sign your immune system is responding appropriately. In most cases, these effects will go away in a few days.

WHEN SHOULD I CALL 911 OR GO TO THE EMERGENCY ROOM?

1 Throat or facial swelling

2 Difficulty breathing or chest pain

3 Stroke-like symptoms: Facial drooping, weakness, confusion

4 Rapid heartbeat or fainting

5 Fever greater than 102F that does not resolve with Tylenol or Ibuprofen

6 Severe rash or hives



FMDH
PUBLIC HEALTH